

# Optical Emission and Parameters of DBD Plasma in a Mixture of Mercury Diiodide and Mercury Dibromide Vapor and Helium with Addition of Xenon

Antonina Malinina\*

Department of Quantum Electronics, Uzhhorod National University, Narodna pl. 3, Uzhhorod, Ukraine

\*Corresponding Author: Antonina Malinina, Department of Quantum Electronics, Uzhhorod National University, Narodna pl. 3, Uzhhorod, Ukraine.

**Abstract:** The spectral, energy, and electrical characteristics of atmospheric pressure nanosecond DBD plasma in HgI<sub>2</sub> / HgBr<sub>2</sub> / Xe / He mixtures were studied. The discharge was initiated by a repetitively pulsed voltage with a pulse repetition rate of 5-20 kHz of positive polarity, an amplitude of 25.5 kV, and a half-height duration of 150 ns. The optical emission of the exciplex molecules HgI(B<sup>2</sup>Σ<sup>+</sup>+1/2-X<sup>2</sup>Σ<sup>+</sup>+1/2) and HgBr(B<sup>2</sup>Σ<sup>+</sup>+1/2-X<sup>2</sup>Σ<sup>+</sup>+1/2) was investigated. Based on a comparison of the temporal behavior of the B-X transition spectra of the HgI (λ<sub>max.</sub>=443 nm) and HgBr (λ<sub>max.</sub>=502 nm) molecules, the mechanism of simultaneous formation of the exciplex sHgI\* and HgBr\* molecules is determined. Specific power radiation, namely 96 W/cm<sup>3</sup> in a single pulse and the average, namely 260 mW/cm<sup>3</sup> at a pulse repetition rate of 18 kHz was achieved. The distribution of the energy and rates of the processes responsible for the emission of HgI and HgBr molecules are analyzed. Studies have revealed the possibility of optimizing the pulsed power in order to obtain simultaneous emission in the visible (violet-blue and blue-green light) spectral range in the plasma of the atmospheric-pressure DBD in the HgI<sub>2</sub>/HgBr<sub>2</sub>/Xe/He mixture.

**Keywords:** DBD plasma, mercury diiodide, mercury dibromide, helium, xenon, optical emission

## 1. INTRODUCTION

The emission characteristics and parameters of the plasma are important both for the improvement of existing and for the creation of new radiation sources [1-3]. Plasma in mixtures of mercury diiodide and mercury dibromide vapor with atomic gases can be an effective source of dual-band coherent and spontaneous emission in the violet-blue and blue-green spectrum with an emission maximum at a wavelength of (λ<sub>max.</sub>) 443 nm and (λ<sub>max.</sub>) 502 nm [4,5]. Light sources that operate on these mixtures provide high spectral radiation power, which is important for practical applications, such as sources of active photosynthetic radiation or lasers and others [6-8]. The high radiation efficiency of such mixtures was demonstrated in the studies of the authors [9-11].

This paper is devoted to the study of optical emission (350-700 nm) of a dielectric barrier discharge (DBD), which was created by a nanosecond (150 ns) pulse-periodic (5-20 kHz) high voltage in a mixture of mercury diiodide, mercury dibromide vapor with helium with the addition of xenon. On the basis of experiments and numerical calculations of plasma parameters, the mechanism of formation of exciplex molecules HgI\* and HgBr\* and additional processes that increase the population of B<sup>2</sup>Σ<sup>+</sup><sub>1/2</sub> -state of mercury monohalides are analyzed.

## 2. TECHNIQUE AND METHODS OF RESEARCH

**The technique and methodology for studying the optical emission of DBD plasma in a mixture of mercury diiodide and mercury dibromide vapor, helium, and xenon was similar to that used in studies of the mixture without the addition of xenon [11].**


The investigated mixture was prepared directly in the interelectrode space. Mercury dihalides: mercury diiodide and mercury dibromide in an amount of 80 mg were preloaded into the interelectrode space. The surfaces of the elements of the internal volume of the tube were outgassed by heating it at a temperature of 50° C and pumping it to a residual gas pressure of 1 Pa for 2 hours. The partial vapor pressure of HgI<sub>2</sub> and HgBr<sub>2</sub> in the mixture was created due to the heating of the mixture during the dissipation of the energy of a repetitively pulsed discharge. It was measured by the

temperature of the coldest part of the tube based on the linear interpolation of the reference data from [12]. The partial pressure of xenon and helium was measured by a sample membrane gauge and pressure gauge, respectively.

### 3. RESEARCH RESULTS AND DISCUSSION


The radiator emission spectra were studied in the region of partial pressures of mercury diiodide and mercury dibromide vapor of 0.1–0.5 kPa, xenon 0.1–4 kPa, and helium 100–180 kPa.

The survey radiation spectrum at a pulse repetition rate of  $f = 18$  kHz is shown in Fig.1. The amplitude of the voltage on the electrodes and the current through the gas-discharge gap is 25.5 kV and 280A, respectively. The total pressure of the mixture is 181.4 kPa. A characteristic feature of this mixture is the presence of systems of spectral bands of the electron-vibrational transition  $B^2\Sigma_{1/2}^+ \rightarrow X^2\Sigma_{1/2}^+$  of the exciplex molecules  $HgI^*$  and  $HgBr^*$  5 and 7 nm wide with radiation maxima at wavelengths  $\lambda = 443$  nm and  $\lambda = 502$  nm, respectively [13], a steep increase in the intensity of these spectral bands on the side of the long-wavelength region and a slow decline in the short-wave region. The edges of the spectral bands overlap the wavelength range of 350–510 nm. In addition to these spectral bands, radiation is also observed on the  $\lambda = 546$  nm line of mercury atoms, the transition  $7s^3S_1 - 6p^3P_2$ , and the  $\lambda = 823$  nm line of xenon atoms, transition  $6s[3/2]_2 - 6p[3/2]_2$  [14,15]. With a change in the pulse repetition frequency from 5000 Hz to 20,000 Hz, the radiation intensity in the spectral bands and lines increases 4 times.




**Fig1.** Survey plasma emission spectrum of a barrier discharge. Partial pressures: mercury diiodide — 0.1 kPa, mercury dibromide — 0.3 kPa, xenon — 1 kPa, helium — 180 kPa. The amplitudes of the voltage and current pulses are 26 kV and 280 A, respectively.

The results of studies of integral characteristics (the dependence of the average radiation power on the partial pressure of the buffer gas helium, xenon, and the time of operation of the emission source in one portion of the working mixture are shown in Fig. 4, Fig. 5 and Fig. 6. The regularity is established with increasing partial pressure of helium from 140 kPa to 200 kPa, an increase in the specific average power, the achievement of its maximum specific value of 260 mW/cm<sup>3</sup> at 180 kPa and a decrease in the further increase in helium pressure is observed.




**Fig2.** The dependence of the specific average radiation power on the partial pressure of helium. The partial pressure of xenon -1 kPa. The pump pulse repetition rate is 18 kHz, the voltage pulse amplitude is 25.5 kV.



**Fig3.** The dependence of the radiation intensity of mercury monohalides in relative units on the pressure of xenon: 1-molecule  $HgI^*$ , 2-molecule  $HgBr^*$ . Helium partial pressure is 180 kPa. The pump pulse repetition rate is 18 kHz, the voltage pulse amplitude is 25.5 kV.


Fig.3 shows the results on the dependences of the radiation intensity of exciplex molecules of mercury monoiodide and mercury monobromide on the partial pressure of xenon. The most intense radiation of these molecules occurs at a partial pressure of xenon in the region of 1 kPa. In the mixture, mercury monoiodide molecules emit more intensively (3.5 times) compared to the radiation intensity of mercury monobromide molecules.

The dependence of the radiation intensity of mercury monohalides in relative units on the number of pump pulses is shown in Fig. 6. The saturation of the radiation power occurs for exciplex molecules  $HgI^*$  and  $HgBr^*$  after  $4 \cdot 10^7$  pulses.



**Fig4.** The dependence of radiation intensity of mercury monohalides on the total number of pulses: 1.  $HgI^*$  molecule, 2.  $HgBr^*$  molecule. The total pressure of the mixture is 181.4 kPa. The pulse repetition rate is 18 kHz, the voltage pulse amplitude is 25.5 kV.

The results of studies of the temporal characteristics of the radiator are shown in Fig.5 with the most optimal radiation power for the experimental conditions. The error and reproducibility of the results of oscillographic measurements were 10% and 90%, respectively. Current pulses (I) have the different polarity. Their complex form is caused by recharging the dielectric-plasma circuit. Their amplitude and duration are  $\sim 280$  A and  $\sim 150$  ns, respectively. Radiation pulses (P) have the same polarity. The pulse durations at half-height were  $\sim 100$  ns and  $\sim 150$  ns for the  $HgI^*$  and  $HgBr^*$  molecules, respectively. The specific power in the radiation pulse is  $96 \text{ W/cm}^3$ .




**Fig5.** Oscillograms of current (I) and radiation (P) pulses: ---  $HgI^*$  molecule, -  $HgBr^*$  molecule. The total pressure of the mixture is 181.4 kPa. The pulse repetition rate is 18 kHz, the voltage pulse amplitude is 25.5 kV.

Two-hump temporal dependence is observed for the emission of both HgI\* and HgBr\* exciplex molecules. The amplitudes of the current and radiation pulses coincide in time within the measurement error (10%). The amplitude of the second radiation pulse for these molecules is higher than the first. The second pulse is characterized by an increase in its duration compared to the first pulse.

The parameters of DBD plasma are optimal for obtaining the maximum radiation power on the mixture HgBr<sub>2</sub>-HgI<sub>2</sub>-Xe-He = 0.001635-0.0005513-0.005514- 0.9823 at a total mixture pressure of 181.4 kPa on the reduced electric field (E/N): were determined numerically and calculated as total integrals of the electron energy distribution function (EEDF) in the discharge. The EEDFs were found numerically by solving the Boltzmann kinetic equation in the two-term approximation [16]. Calculations of the EEDF were performed using the program [17]. Based on the resulting EEDF, mean electron energy, the specific power losses of the electric discharge on various elementary processes in the plasma, and the rate constants of elastic and inelastic scattering of electrons on mercury diiodide and mercury dibromide molecules, xenon and helium atoms depending on the value of the reduced electric field (ratio electric field strength (E) to the total concentration of mercury dihalide molecules, xenon and helium atoms (N)). The range of changes in the parameter E/N=1-100 Td (1·10<sup>-17</sup> - 1·10<sup>-15</sup> V·cm<sup>2</sup>) and included the values of the parameter E/N, which were implemented in the experiment.


The integral of electron collisions with helium, xenon, mercury dibromide and mercury diiodide molecules includes the following processes of elastic and inelastic collisions: elastic scattering, excitation of energy states of the atom He (2<sup>3</sup>S, 2<sup>1</sup>S, 2<sup>3</sup>P, 2<sup>1</sup>P, 3SPD, 4SPD, 5SPD), helium atom ionization, elastic scattering and excitation of electronic states of xenon atom with threshold energies: 3.4 eV, 8.31 eV, 8.44 eV, 9.69 eV, 10.0 eV, 11.0 eV, 11.7 eV, ionization of xenon atoms, vibrational excitation of HgBr<sub>2</sub> molecules, excitation (HgBr<sub>2</sub> (D)) - state of HgBr<sub>2</sub> molecules, dissociative excitation of mercury monobromide(X<sup>2</sup>Σ<sup>+</sup><sub>1/2</sub>, B<sup>2</sup>Σ<sup>+</sup><sub>1/2</sub>- states), attachment, ionization of mercury dibromide molecules, dissociative excitation of B<sup>2</sup>Σ<sup>+</sup><sub>1/2</sub>-state of mercury monoiodide and ionization of mercury diiodide molecules. The data on the absolute values of the effective cross sections for these processes, as well as their dependence on the electron energy, are taken from [17, 18–21].

Fig.6 shows the characteristic view of the EEDF when changing the reduced electric field strength (E / N) in the range of 1-100 Td. The parameters have been determined on dependence on the reduced electric field (E / N) in the range of 1-100 Td. An increase of its value leads to increase of the number of "fast" electrons in the discharge. Mean electron energy of the discharge increases sharply from 0.5 to 3.8 eV with increasing E/N from 1 to 14.7 Td. With further increase of the reduced electric field the rate of increase of mean electron energy slows down and reaches a value of 12.2 eV for E/N = 100 Td. The value of the reduced electric field at which experimental studies were carried out and the mean electron energy and their temperature is equal to 68 Td, 9.6 eV and 111360K, respectively. The concentration of electrons was 4.6 · 10<sup>17</sup> m<sup>-3</sup>.




**Fig6.** The electron energy distribution functions in the mixture: HgBr<sub>2</sub>: HgI<sub>2</sub>:Xe: He=0.00163 5:0.0005513:0.005514:0.9823 with a total mixture pressure P = 181.4 kPa for the values of the parameter E/N: 1 (1), 25.8 (2), 50.5 (3), 75.2 (4), 100 (5) Td; The inset shows the dependence of mean electron energy on the parameter E/N.

Fig.7 shows the results of studies of the dependence of the rate constants of inelastic collisions of electrons with mercury dihalide molecules on the parameter E/N. For a reduced electric field value of 68 Td, at which experimental studies were performed, the rate constants of dissociative excitation by electrons of the  $B^2\Sigma^+_{1/2}$ -state of mercury monoiodide (1) and mercury monobromide (2) molecules are  $2 \times 10^{-14} \text{ m}^3 / \text{s}$ .



**Fig7.** The dependence of the collision rate constants of electrons with components of a gas-discharge plasma on a mixture of  $\text{HgBr}_2 : \text{HgI}_2 : \text{Xe} : \text{He} = 0.001635:0.0005513:0.005514:0.9823$  with a total mixture pressure  $P = 181.4 \text{ kPa}$  on the parameter E/N: 1 - dissociative excitation of  $B^2\Sigma^+_{1/2}$ -states of the  $\text{HgI}^*$  molecules, 2 - dissociative excitation of  $B^2\Sigma^+_{1/2}$ -states of  $\text{HgBr}^*$  molecules; 3 - dissociative excitation of the  $X^2\Sigma^+_{1/2}$ -state of  $\text{HgBr}$  molecules; 4 - the excitation of ( $\text{HgBr}_2(D)$ ) states of  $\text{HgBr}_2$  molecules.

The distribution of the specific power losses of the discharge on the main processes is shown in Fig. 10. The fraction of the specific power of the discharge which goes on the processes of dissociative excitation of mercury monobromide and mercury monoiodideexciplex molecules increases with the increase of the parameter E/N. It reaches a maximum of 10.6%, 25.5%, 35.7% and 3.3% with the E/N parameter equal to 21.5 Td, 18.1 Td, 31.7 Td and 18.1 Td for electronic states of mercury monobromide  $B^2\Sigma^+_{1/2}$ ,  $X^2\Sigma^+_{1/2}$ , mercury dibromide ( $\text{HgBr}_2(D)$ ) and  $B^2\Sigma^+_{1/2}$ -state of mercury monoiodide, respectively, and with a further increase of parameter E/N it decreases.






**Fig8.** The dependence of the specific power losses of the discharge on the processes of electroncollisionson the  $\text{HgBr}_2 : \text{HgI}_2 : \text{Xe} : \text{He} = 0.001635:0.0005513:0.005514:0.9823$  mixture at the total mixture pressure( $P = 181.4 \text{ kPa}$ ) on the E/Nparameter: 1 -the dissociative excitation of  $B^2\Sigma^+_{1/2}$ -state of  $\text{HgI}^*$  molecules, 2 –the dissociative excitations of  $B^2\Sigma^+_{1/2}$ -state of  $\text{HgBr}^*$  molecules, 3 – the dissociative excitations of  $X^2\Sigma^+_{1/2}$ -state of  $\text{HgBr}^*$  molecules, 4-excitation of the electronicstateofmercurydibromide ( $\text{HgBr}_2(D)$ ).

The emission of spectral bands and lines is probably observed due to the following reactions [16,18-24]:







$$\lambda_{\text{max.}} = 443 \text{ nm}, \lambda_{\text{max.}} = 502 \text{ nm} \quad (6)$$



$$\lambda = 546 \text{ nm},$$



$$\lambda = 823 \text{ nm},$$

where X is a halogen atom (I, Br), M is the concentration of quenching molecules and atoms (HgX<sub>2</sub>, Xe, He), respectively, ΔE<sub>1,2</sub> is the energy difference in reactions (4).

Reactions (1) and (2) are the main sources of formation of exciplex molecules HgX\* [18 - 23]. Their rate constant has a value  $2 \times 10^{-14} \text{ m}^3/\text{s}$  (Fig.7) for our experimental conditions with a reduced electric field strength of 68 Td.

In addition, the exciplex molecules of mercury monoiodide and mercury monobromide are formed due to reactions (3, 4) due to the excitation of mercury diiodide and mercury dibromide molecules by electrons to state D [20]. The rate constant has a value of  $9.9 \times 10^{-14} \text{ m}^3/\text{s}$  for mercury dibromide molecules (Fig.7). This state of mercury dihalide molecules is the sum of all states that are located between the threshold energy and the ionization energy. Emission from the D states of the HgX<sub>2</sub> molecules is not observed, due to the fact that these states dissociate with the formation of mercury monohalides in the (C, D) states. In our experimental conditions, emission from C and D states of mercury monoiodide and mercury monobromide molecules is not observed due to the high efficiency of the quenching process (4) (It is known that the product of the quenching rate constant by helium and xenon of C<sup>2</sup>Π<sub>1/2</sub>-state of mercury monoiodide molecules on the lifetime is equal to  $4.2 \cdot 10^{-18} \text{ cm}^3$  and  $4.2 \cdot 10^{-18} \text{ cm}^3$ , respectively, and the process of quenching (C, D)-states by mercury dihalides occurs more efficiently [24, 25]). The population of these states is transferred to B<sup>2</sup>Σ<sup>+</sup><sub>1/2</sub> - state of mercury monoiodide and mercury monobromide molecules or to other non-optical channels [25]. The reaction of the collision of mercury dihalide molecules with electrons (5) is the channel for the formation of mercury monohalides molecules in the ground state, the rate constant of which is  $3.9 \cdot 10^{-14} \text{ m}^3/\text{s}$  (Fig.7). Electron-vibrational transitions B<sup>2</sup>Σ<sup>+</sup><sub>1/2</sub> → X<sup>2</sup>Σ<sup>+</sup><sub>1/2</sub> of mercury monoiodide and mercury monobromide molecules lead to emission of spectral bands with a maximum intensity at wavelengths λ<sub>max.</sub> = 443 nm and λ<sub>max.</sub> = 502 nm (reaction 6). In the quenching reaction (7), an electron-vibrational transition of mercury dihalide molecules to their ground state occurs without radiation. The rate constant of this process for quenching of mercury monoiodide and mercury monobromide molecules by helium <math>2.9 \cdot 10^{-20} \text{ m}^3/\text{s}</math> and <math>3.4 \cdot 10^{-20} \text{ m}^3/\text{s}</math>, respectively [25] and by mercury dihalides  $1.1 \cdot 10^{-16} \text{ m}^3/\text{s}$  and  $3 \cdot 10^{-16} \text{ m}^3/\text{s}$  [24].

Excited mercury atoms are formed due to the passage of reaction (8) due to the large effective cross section of the dissociative excitation of HgX<sub>2</sub> molecules. Their emission (reaction 9) leads to the appearance of the line λ = 546nm (Fig.1). Reaction (10) is responsible for the excitation of xenon atoms, the rate constant, which is  $\sim 1 \cdot 10^{-17} \text{ m}^3/\text{s}$  (our calculation data). Excited xenon atoms emit quanta (hν) as a result of radiative decay as a result of which a xenon line λ=823 nm was observed in the spectrum (Fig. 2) (reaction 10). In addition, the presence in the emission spectra (Fig. 2) of the line of Hg atoms can be caused by the process of degradation of the working mixture. Part of mercury monoiodide and mercury monobromide molecules does not have time to associate in the processes:





dissociate into mercury and halogen atoms in collisions with electrons [26]. Halogen atoms have time to escape to the surface of the electrode and form metal halides [27,28]. Mercury atoms accumulate and, in collisions with electrons, are excited and emit radiation  $\lambda = 546$  nm.

A steep increase in intensity from the part of the spectrum with large wavelengths and a slow decline in the short-wavelength region (Fig.1) for both the mercury monoiodide molecule and mercury monobromide molecule are explained by the form of potential curves (excited  $B^2\Sigma^+_{1/2}$  – state is shifted towards large inter nuclear distances relative to  $X^2\Sigma^+_{1/2}$  - state) and relaxation processes of the population of the upper vibrational levels of the excited electronic state, which occur faster than the electronic-vibrational transition to the main  $X^2\Sigma^+_{1/2}$ -state [29,30].

The regularity of the increased intensity of  $\text{HgI}^*$  molecules compared to the intensity of  $\text{HgBr}^*$  molecules at the same excitation rate constants ( $2.0 \times 10^{-14}$  m<sup>3</sup>/s, Fig.7) is caused by the quenching process of the  $B^2\Sigma^+_{1/2}$  - state of mercury monohalides by mercury dihalide molecules (reaction 7). The quenching rate constant of the  $B^2\Sigma^+_{1/2}$ -state of mercury monoiodide molecules - is  $1.1 \cdot 10^{-10}$  cm<sup>3</sup>/s, and that of mercury monobromide molecules –  $3.4 \cdot 10^{-10}$  cm<sup>3</sup>/s [24], which makes the difference in the emission intensities of these molecules.

The dependence of the radiation power on the partial pressure of helium (Fig.2) is caused primarily by the following processes: an increase in the electron concentration with an increase in the partial pressure of helium in the mixture, a change in the fraction of the discharge energy spent on heating the working mixture; a change in mean electron energy and the rate constant of the excitation of the exciplex molecules  $\text{HgI}^*$  and  $\text{HgBr}^*$  depending on the values of the parameter E/N, as well as quenching process of the  $B^2\Sigma^+_{1/2}$  – state of molecules  $\text{HgI}^*$  and  $\text{HgBr}^*$  in collisions with helium atoms, xenon and mercury dihalide molecules. With an increase in the partial pressure of helium in the mixture, the value of the parameter E/N decreases. This leads to an increase in the specific losses of the discharge power on the elastic scattering of electrons on atoms and molecules (Fig.8) and, accordingly, to an increase in the partial pressure of mercury diiodide and mercury dibromide vapor and the radiation power of mercury monoiodide and mercury monobromide exciplex molecules. In addition, an increase in the electron concentration increases with increasing partial pressures of both helium and mercury diiodide and mercury dibromide vapor, also increases the concentration of electrons, which increases with increasing concentration of the components of the working mixture [1]. The presence of a maximum and a further drop in the radiation power of the exciplex molecules  $\text{HgI}^*$  and  $\text{HgBr}^*$  with an increase in the partial pressure of helium is associated with a decrease in the fraction of the discharge energy that is spent on exciting their  $B^2\Sigma^+_{1/2}$  – state (Fig.8), as well as well as quenching the states of mercury monohalide molecules when they collide with helium atoms [1].



Where  $\Delta E_{3,4}$  is the energy difference in the reactions.

The dependence of the intensity of the emission of  $\text{HgI}^*$  and  $\text{HgBr}^*$  molecules on the partial pressure of xenon (Fig. 5) is determined by the efficiency of the processes leading to their excitation and quenching —  $B^2\Sigma^+_{1/2}$ -states of these molecules in the DBD plasma. The quenching rate constant for xenon of  $\text{HgI}^*$  and  $\text{HgBr}^*$  molecules is  $2.2 \times 10^{-13}$  cm<sup>3</sup>/s и  $3.8 \times 10^{-14}$  cm<sup>3</sup>/s, respectively [25].

The earlier achievement of saturation in the intensity of  $\text{HgBr}^*$  radiation from the number of pulses than for  $\text{HgI}^*$  molecules (Fig. 4) is caused by a greater loss of discharge power on the dissociative excitation of  $B^2\Sigma^+_{1/2}$  – state of  $\text{HgBr}^*$  molecules compared to the  $\text{HgI}^*$  molecules (Fig.8).

The temporal dependence of the current pulses (Fig.5) is caused by the charge exchange of the dielectric-plasma circuit. The increased amplitudes of the second radiation pulse, its duration and the duration of the falling front, as compared with the first pulse, are caused by the increase in the population of the  $B^2\Sigma^+_{1/2}$ - state of the  $\text{HgI}^*$  and  $\text{HgBr}^*$  molecules, which pass on time later than the main (dissociative excitation by electrons in collisions with mercury dihalide molecules). Such processes can be:



The process of recombination of positive mercury dihalide ions with electrons (17), according to the authors of [31], can play a significant role at current densities of  $\sim 1000 \text{ A/cm}^2$ , which is not observed under the conditions of our experiment. The calculated rate constants of the recombination process (18) of a positive ion with a negative ion are high (for  $\text{HgI}_2$  and  $\text{HgBr}_2$  molecules  $\sim 3 \times 10^{-7} \text{ cm}^3/\text{s}$ ), however, under gas-discharge conditions due to low concentrations of atomic ion (rate constant of the process of negative ions formation  $\text{I}^- \text{ Br}^- \sim 10^{-10} \text{ cm}^3/\text{s}$  [21]) its role is negligible. The excitation of radicals  $\text{HgX}(\text{X}^2\Sigma_{1/2}^+)$  by electrons in the  $\text{B}^2\Sigma_{1/2}^+$ -state (19) is possible under our experimental conditions.

The regularity of the difference in the magnitudes of the amplitudes of the first and second radiation pulses (Fig.5) is explained as follows. The first and second excitation pulses (current) form mercury monoiodide and mercury monobromide molecules in  $\text{B}^2\Sigma_{1/2}^+$  and  $\text{X}^2\Sigma_{1/2}^+$ -states due to dissociation of mercury dihalide molecules in collisions with electrons. The second excitation pulse (current), in addition, leads to an additional increase in the population of the  $\text{B}^2\Sigma_{1/2}^+$ -state of mercury monobromide molecules due to the process (19), where  $\text{HgX}(\text{X}^2\Sigma_{1/2}^+)$  mercury monoiodide and mercury monobromide molecules in the ground state and that did not have time to recover in triatomic molecules (mercury diiodide and mercury dibromide) in the interpulse period (150 ns) in the processes (13,14), which leads to an increase in the amplitude of the second radiation pulse and an increase in its duration compared to the first pulse.

Based on the calculated reaction rate constants (Fig.7) and specific power loss distribution (Fig.8) as a function of  $E/N$ , appears the possibility to assume the optimal power input scheme in order to get the maximum possible emission from simultaneous violet - blue ( $\lambda=444 \text{ nm}$ ) and blue-green ( $\lambda=502 \text{ nm}$ ) spectral ranges for the  $\text{HgI}_2 / \text{HgBr}_2/\text{Xe} / \text{Hemixture}$ . Indeed, the excitation of the levels of the main reactions is attained at 11-30 Td (see Fig.8).

This field is too small for effective ionization of the mixture and stable development of the discharge. At pressures higher than 1 Atm, a much higher electric field is required, at least 100 Td. Thus, a mixed power source, when a short powerful pulse of a few nano seconds in duration (3-5) is used to obtain a relatively high initial ionization ( $10^{12}-10^{13} \text{ cm}^{-3}$ ), and then a long pulse of ten hundreds of nano seconds, depending on the component composition of the mixture and pressure with relatively low electric fields, provide effective excitation of  $\text{HgI}^*$  and  $\text{HgBr}^*$  molecules, can be the optimal solution for discharge lamps based on  $\text{HgI}^*$  and  $\text{HgBr}^*$  emission.

#### 4. CONCLUSION

The simultaneous emission from two exciplex molecular bands  $\text{HgI}^*$  ( $\lambda=444 \text{ nm}$ ) and  $\text{HgBr}^*$  ( $\lambda=502 \text{ nm}$ ) in a nano second DBD in a mixture of  $\text{HgI}_2/\text{HgBr}_2/\text{Xe}/\text{He} = 0.001635/0.0005513/0.005514/0.9823$  at atmosphere pressure is studied. Specific power radiation in a single pulse-  $96 \text{ W/cm}^3$  and the average-  $260 \text{ mW/cm}^3$  at a pulse repetition rate of 18 kHz is achieved. Long-term stable more than  $7 \cdot 10^7$  pulses (within a measurement error of 10%) emission was detected.

The current and radiation shape of  $\text{HgI}^*$  and  $\text{HgBr}^*$  molecules with a temporal resolution are investigated. It is established that  $\text{HgI}^*$  radiation repeats the form of  $\text{HgBr}^*$  radiation and is close in the time scale.

Based on the analysis of possible reactions of excitation and quenching, it was found that the population of the exciplex molecules  $\text{HgI}^*$  and  $\text{HgBr}^*$  ( $\text{B}^2\Sigma_{1/2}^+$ )-state takes place mainly due to dissociative excitation in collisions with electrons. A discussion of additional processes that increase the population of the upper  $\text{B}^2\Sigma_{1/2}^+$ -state of mercury monohalides is given.

As a result of calculating the reaction rates and energy distribution for different electric fields, it can be concluded that a combination of a high electric field at the front of the discharge and a low field over the duration of the pulse can be used to increase the emission power in the mixture studied.



Based on the results of plasma emission investigations of a barrier discharge in a mixture of mercury diiodide and mercury dibromide vapor with xenon and helium, an efficient source of simultaneous two-band emission in the violet-blue and blue-green spectral ranges can be created, which will be used as a source of active photosynthetic radiation and a laser in the field of biotechnology, photonics, medicine, etc.

#### ACKNOWLEDGMENTS

The author is grateful to prof. Alexander Malinin and prof. Alexander Shuaibov for help in discussing the results.

#### REFERENCES

- [1] Raizer, Yu.P., Gas Discharge Physics, (Moscow: Nauka, 1991), p. 449.
- [2] McDaniel, E. W. and Nigan, W.L., Gas Lasers, (New York: Academic Press, 1982), p. 486.
- [3] Kogelschatz, U., Ultraviolet excimer radiation from nonequilibrium gas discharges and its application in photophysics, photochemistry and photobiology. J. Opt. Technol. 79, 484 (2012).
- [4] Berry, A.J., Whitehurst, C. and King, T. A., Multihalide operation of mercury halide lasers. J. Phys. D. Appl. Phys., 21, 39 (1988).
- [5] Bazhulin, S. P., Basov, N. G., Bugrimov, S. N., Zuev, V. S., Kamrukov, A. S., Kashnikov, G. N., Kozlov, N. P., Ovchinnikov, P. A., Opekan, A. G., Orlov, V. K. and Protasov, Yu. S., Blue-violet HgI/HgI<sub>2</sub> laser with wide-band optical pumping by a linearly stabilized surface discharge. Kvantovaya Elektronika, 13(5), 1017 (1986).
- [6] Posudin, Yu.I., Laser Photobiology (Kiev: Vishchashkola, 1989), p.248.
- [7] Romanenko, V.D., Krot, Yu.G., Sirenko, L.A. and Solomatina, V.D., Biotechnology of Hydrobion Cultivation' (Kiev: Institute of Hydrobiology, NAS Ukraine, 1999), p. 264.
- [8] Sosnin, Edward A., Oppenlander, Thomas, Tarasenko, Victor F., Applications of capacitive and barrier discharge excilamps in photoscience. Journal of Photochemistry and Photobiology C: Photochemistry Reviews, 7, 145 (2006).
- [9] Malinin, A. N., Optical characteristics of the gas-discharge plasma of working mixtures in an excimer HgBr/HgI radiation source', Kvantovaya Elektronika. 35(3), 243 (2005).
- [10] Datsyuk, V.V., Izmailov, I.A., Naumov, V.V., Kochelap, V.A., Direct electron-impact mechanism of excitation of mercury monobromide in a double-pulse dielectric-barrier-discharge HgBr lamp. Plasma Sources Sci. Technol. 25, 045020 (2016).
- [11] Malinina, A.A., Malinin, A.N. and Shuaibov, A.K., Optical Emission and Parameters of Dielectric Barrier Discharge Plasma In A Mixture of Mercury Diiodide and Mercury Dibromide Vapor With Helium. Open Physics Journal. 4, 43 (2018).
- [12] Efimov, A.E., Belorukova, L.P., Vasilkova, I.V., Chechev, V.P., Properties of inorganic compounds. Reference book (L: Chemistry, 1983), p. 392.
- [13] Pears, R.W., Gaydon, A.G., The identification of molecular spectra (Ldn Chopman Holl LTD, 1963), p.347.
- [14] Zaydel, A.N., Prokofiev, V.K., Raysky, S.M., Slavny V.A. and Shreider, E.Ya., Tables of spectral lines (M.: Nauka, 1977), 800p.
- [15] Prokopiev, V.E., Yatsenko, A.S., Energy levels and radiative transitions of neutral atoms, (Novosibirsk: Preprint / IAE SB AS USSR, No. 161, 1981), p. 52.
- [16] Hagelaar, G. J. M. and Pitchford, L. C., Solving the Boltzmann equation to obtain electron transport coefficients and rate coefficients for fluid models. Plasma Sources Sci. Technol. 14, 722 (2005).
- [17] <http://www.bolsig.laplace.univ-tlse.fr/>
- [18] Kushawaha, V. and Mahmood, M., Electron impact dissociation of HgX<sub>2</sub> (X=Cl, Br, I). J. Appl. Phys., 62(6), 2173 (1987).
- [19] Malinin, A.N., Excitation of the B<sup>2</sup>Σ<sub>1/2</sub><sup>+</sup> state of mercury monohalides by electron impact, Laser Physics, 7(6), 1168 (1997).
- [20] Nighan, W.L. and Brown, R.T., Kinetic processes in the HgBr(B → X) HgBr dissociation laser. J. Appl. Phys., 53(11), 7201 (1982).
- [21] Malinin, A. N. and Shimon, L. L., Excitation of the B<sup>2</sup>Σ<sub>1/2</sub><sup>+</sup> state of the HgBr\* molecules in a gas-discharge plasma formed from a mixture of mercury dibromide and helium. Kvantovaya Elektronika, 23(12), 1077 (1996).

- [22] Malinin, A.N., Shuaibov, A.K. and Shevera, V.S., Dissociative excitation of the  $B^2\Sigma^+_{1/2}$  states of mercury monohalide by electron impact. Soviet Journal of Quantum Electronics, 13(7), 977 (1983).
- [23] Malinin, A.N. The Main Characteristics of the Plasma of Pulse Glow Discharge in Mixtures of Mercury Dihalide and Rare Gases. Laser Physics, 8(2), 395 (1998).
- [24] Malinina A.A., Malinin A.N. Efficiency of quenching of the  $B^2\Sigma^+_{1/2}$  state of mercury monoiodide and monobromide by mercury dihalides in gas-discharge plasma in multicomponent mixtures of an exciplex radiation source. Optics and Spectroscopy, 105(1), 32 (2008).
- [25] Roxlo, C. and Mandl, A., Quenching kinetics for the HgBr ( $B^2\Sigma_{1/2}$ ) and HgJ ( $B^2\Sigma_{1/2}$ ,  $C^2\Pi_{1/2}$ ) states. J.Chem.Phys., 72(1), 541 (1980).
- [26] McDaniel, E., and Nighan, W., Gas Lasers, Eds. (New York: Academic, 1982).
- [27] Erlason, A.C., Cool, T.A., On the regeneration mechanism of HgBr<sub>2</sub> in HgBr/HgBr<sub>2</sub> dissociation lasers. ChemPhysLett., 96, 685 (1983).
- [28] Liu, C.S., Liberman, Irving, Chemical Control of HgBr Lasers. Quantum Electronics, 23(2), 245 (1987).
- [29] Wadt, W.R., The electronic structure of HgCl and HgBr. Appl. Phys. Lett, 34, 658 (1979).
- [30] Datsyuk, V.V., Izmailov, I.A. and Kochelap, V.A., Vibrational relaxation of excimers. Physics Uspekhi, 41, 379 (1998).
- [31] Burhman, R, Schimitschek, E.J., High-power blue-green lasers. Laser Focus, 6, 54 (1981).

**Citation:** Antonina Malinina. " Optical Emission and Parameters of DBD Plasma in a Mixture of Mercury Diiodide and Mercury Dibromide Vapor and Helium with Addition of Xenon", *International Journal of Advanced Research in Physical Science (IJARPS)*, vol. 5, no. 12, pp. 10-19, 2018.

**Copyright:** © 2018 Authors, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.