

In Silico Phylogenetic Study of the Human Tp53 Gene: Insights into Evolutionary Dynamics

Akhil Kumar Pigili¹, Anil Kumar Babu Katika²

¹Department of Bioinformatics, Sri Venkateswara Institute of Medical Sciences, Tirupati, India

²Department of Biotechnology, Central Tribal University of Andhra Pradesh, Vizianagaram, India

***Corresponding Author:** Akhil Kumar Pigili, Department of Bioinformatics, Sri Venkateswara Institute of Medical Sciences, Tirupati, India.

Abstract

The tumour suppressor protein p53, sometimes known as the "protector of the genome," is widely recognised for its crucial function in averting cancer through the regulation of DNA repair, disruption of the cell cycle, and programmed cell death known as apoptosis. Recent research has shown that p53 plays a more complex role in genomic equilibrium than its conventional roles. The Tp53 gene's evolutionary linkages are examined in this work while evolutionary studies are crucial to contemporary scientific research. Studies on evolution provide links that are useful for current study. The *Homo sapiens* tumour protein Tp53 transcript variant-1 mRNA FASTA sequence used in this work was obtained from the NCBI, after the Multiple Sequence Alignment using UPGMA and MEGA were used to conduct evolutionary experiments with other species. Consequently, this research. Thus, this work will aid current biological research in developing and comprehending the evolutionary dynamics and pathways of the Tp53 gene in Clinical Research.

Keywords: p53, tumour suppressor gene, phylogenetics, evolutionary analysis, sequence alignment, BLAST

1. INTRODUCTION

The p53 gene, also known as TP53 or tumour protein, encodes a protein that controls the cell cycle and thereby inhibits cancer formation. In species with many cell lines, the capacity of cells to suppress cancer is essential. P53's role in preserving rigidity by preventing genome changes has given it the nickname "the guardian of the genome" (Strachan and Read, 1999). It has an atomic mass of 53 kilo Dalton, which is typical for cell proteins. It is located on the final region of the human seventeenth chromosome. The structure of p53 includes the p53 protein is made up of phosphoproteins that include 393 amino acids. The tumour silencer p53 is well-known for its role in cell cycle capture, apoptosis, senescence, and Ferro ptosis. As of late, more evidence has emerged suggesting p53 is also effectively involved in the reconstruction of the cellular digestive system. The quantity of scientific evidence supporting p53's multiple activities in metabolism, normal tissue functioning, and tumour progression is expanding at an incredible rate. Owing to the significance of this gene, further research into the regulation and function of p53 is still required in order to provide more effective treatments and diagnostics. Understanding the trends and mechanisms of evolution is crucial for current scientific study, and this gene's phylogenetic analysis plays a key role in that regard. It has shown to be a crucial component in bringing together a wide variety of scientific disciplines to analyse and explore evolutionary patterns. The distribution pattern of p53 genetic variation, which is present in several creatures including humans, is also investigated in this work.

2. MATERIALS AND METHODS

2.1. Retrieval of Nucleotide Sequences of P53 Gene

The nucleotide sequence of *Homo sapiens* tumour protein p53 (Tp53) transcript variant-1 mRNA was retrieved from the NCBI website (<https://www.ncbi.nlm.nih.gov>) with the accession number of gene bank in FASTA format.

2.2. Determination of Identity and Similarity (%)

2.2.1. Local Sequence Alignment

The BLAST (Basic Local Alignment Search Tool) technique was utilized to evaluate the identification and similarity (%) of the chosen animal. Homo sapiens cancer protein Tp53 transcript variant-1 mRNA was obtained from the NCBI website in the FASTA format, and the BLAST N (Nucleotide Blast) was carried out from the NCBI BLAST site (<https://blast.ncbi.nlm.nih.gov>) website. BLAST (Altschul et al., 1990) was carried out for the Tp53 gene.

2.2.2. Phylogenetic Analysis

The CLUSTALW tool was used to align the Tp53 gene sequences multiple and pairwise. Using the web program genome.jp, a phylogenetic analysis of the H. sapiens tumour protein p53 (TP53) and transcript variant-1 mRNA sequence was performed using CLUSTAL W. The program created a phylogenetic tree that illustrated the ancestral connection between the sequences. The 1000 replications of the bootstrap method were used to recreate the evolutionary tree. Sequences that are located in the same cluster have a close relationship. The tree displays many groupings that indicate how they relate to one another.

3. RESULTS AND DISCUSSION

3.1. Sequence retrieval

H. sapiens tumour protein p53 (TP53), transcript variant-1 mRNA was retrieved from the NCBI in FASTA format. The sequence of the transcript variant-1 mRNA (NM_000546.4) is as the following:

>NC_000017.11:c7687490-7668421 Homo sapiens chromosome 17, GRCh38.p14 Primary Assembly

```
GAGCTGAATGAGGCCTTGGAACTCAAGGATGCCAG      TAGGTAGAGGGAGTTGTCAAGTCTCTGCTGGCCCAG
TAGGTAGAGGGAGTTGTCAAGTCTCTGCTGGCCCAG      CCAAACCCTGTCTGACAACCTCTTGGTGAACCTTAGT
CCAAACCCTGTCTGACAACCTCTTGGTGAACCTTAGT      ACCTAAAAGGAAATCTCACCCATCCCACACCCTGG
ACCTAAAAGGAAATCTCACCCATCCCACACCCTGG      AGGATTCATCTCTGTATATGATGATCTGGATCCAC
AGGATTCATCTCTGTATATGATGATCTGGATCCAC      CAAGACTGTTTTATGCTCAGGGTCAATTTCTTTTTTC
CAAGACTGTTTTATGCTCAGGGTCAATTTCTTTTTTC      TTTTTTTTTTTTTTTTTCTTTTTCTTGAGACTGGGTC
TTTTTTTTTTTTTTTTCTTTTTCTTGAGACTGGGTC      TCGCTTGTTGCCAGGCTGGAGTGGAGTGGCGTGA
TCGCTTGTTGCCAGGCTGGAGTGGAGTGGCGTGA      TCTTGGCTTACTGCAGCCTTGCCTCCCCGGCTCGA
TCTTGGCTTACTGCAGCCTTGCCTCCCCGGCTCGA      GCAGTCCTGCCTCAGCCTCCGGAGTAGCTGGGACC
GCAGTCCTGCCTCAGCCTCCGGAGTAGCTGGGACC      ACAGGTTTCATGCCACCATGGCCAGCCAACCTTTGCA
ACAGGTTTCATGCCACCATGGCCAGCCAACCTTTGCA      TGTTTTGTAGAGATGGGGTCTCACAGTGTGCCAG
TGTTTTGTAGAGATGGGGTCTCACAGTGTGCCAG      GCTGGTCTCAAACCTCTGGGCTCAGGCGATCCACCT
GCTGGTCTCAAACCTCTGGGCTCAGGCGATCCACCT      GTCTCAGCCTCCAGAGTCTGGGATTACAATTGTG
GTCTCAGCCTCCAGAGTCTGGGATTACAATTGTG      AGCCACCACGTCCAGCTGGAAGGGTCAACATCTTTT
AGCCACCACGTCCAGCTGGAAGGGTCAACATCTTTT      ACATTCTGCAAGCACATCTGCATTTTCACCCCACCT
ACATTCTGCAAGCACATCTGCATTTTCACCCCACCT      TCCCCTCCTTCCCTTTTTATATCCATTTTATATC
TCCCCTCCTTCCCTTTTTATATCCATTTTATATC      GATCTCTATTTTACAATAAACTTTGCTGCCACCTGT
GATCTCTATTTTACAATAAACTTTGCTGCCACCTGT      GTGTCTGAGGGGT
```

3.2. Local Sequence Alignment

Homo sapiens cancer protein Tp53 transcript variant-1 mRNA was obtained from the NCBI website in the FASTA format, and the BLAST N (Nucleotide Blast) was carried out from the NCBI BLAST site (<https://blast.ncbi.nlm.nih.gov>) website. BLAST (Altschul et al., 1990) was carried out for the Tp53 gene and the results of the BLAST were formulated in a table (table.1) showcasing the E-value, Accession Number, Percent Identity and organism and Sequence details




Figure 1. Genomic Context of Tp53 Gene

Table 1. BLAST Table of Homo sapiens cancer protein Tp53 transcript variant-1 mRNA Gene (Performed Analysis Using NCBI BLAST Program)

S. No	AC Number	E-Value	Organism	Description	Seq Length	PW Identity
1	NM_000546.6	0	Homo sapiens	Homo sapiens tumour protein p53 (TP53), transcript variant 1, mRNA	2512	100
2	HG975427.1	0	Homo sapiens	TPA: Homo sapiens Processed transcript p53-mRNA (p53 gene)	2586	100
3	NM_001276761.3	0	Homo sapiens	Homo sapiens tumour protein p53 (TP53), transcript variant 2, mRNA	2509	99.88
4	AF307851.1	0	Homo sapiens	Homo sapiens p53 protein mRNA, complete cds	2521	99.88
5	BC003596.1	0	Homo sapiens	Homo sapiens tumour protein p53, mRNA (cDNA clone MGC:646 IMAGE:3544714), complete cds	2508	99.88
6	XM_001172077.6	0	Pan troglodytes	PREDICTED: Pan troglodytes tumour protein p53 (TP53), transcript variant X2, mRNA	2537	99.36
7	XM_004058511.5	0	Gorilla gorilla gorilla	PREDICTED: Gorilla gorilla gorilla tumour protein p53 (TP53), transcript variant X2, mRNA	2529	99.36
8	XM_003810066.5	0	Pan paniscus	PREDICTED: Pan paniscus tumour protein p53 (TP53), transcript variant X2, mRNA	2555	99.2
9	DQ191317.1	0	Homo sapiens	Homo sapiens p53 protein (TP53) mRNA, complete cds, alternatively spliced	2584	99.09
10	FJ207420.1	0	Homo sapiens	Homo sapiens mutant p53 mRNA, complete cds	2513	99.09
11	DQ286964.1	0	Homo sapiens	Homo sapiens p53 protein (TP53) mRNA, complete cds, alternatively spliced	2529	99.09
12	NM_001407262.1	0	Homo sapiens	Homo sapiens tumour protein p53 (TP53), transcript variant 9, mRNA	2615	100
13	JN900492.1	0	Homo sapiens	Homo sapiens tumour suppressor TP53 (TP53) mRNA, complete cds, alternatively spliced	2697	100
14	NM_001126118.2	0	Homo sapiens	Homo sapiens tumour protein p53 (TP53), transcript variant 8, mRNA	2629	100
15	XM_016931470.3	0	Pan troglodytes	PREDICTED: Pan troglodytes tumour protein p53 (TP53), transcript variant X1, mRNA	2634	99.3
16	XM_057300362.2	0	Pan paniscus	PREDICTED: Pan paniscus tumour protein p53 (TP53), transcript variant X1, mRNA	2623	99.13
17	NM_001407269.1	0	Homo sapiens	Homo sapiens tumour protein p53 (TP53), transcript variant 12, mRNA	2642	100
18	NM_001276696.3	0	Homo sapiens	Homo sapiens tumour protein p53 (TP53), transcript variant 3, mRNA	2645	100
19	NM_001276695.3	0	Homo sapiens	Homo sapiens tumour protein p53 (TP53), transcript variant 4, mRNA	2572	100

3.3. Phylogenetic Analysis

Finding evidence of directional selection in molecular evolution is a common step in phylogenetic research (Koref et al., 2003; Hsu et al., 2005; Hofmann et al., 2003; Yang and Bielawski, 2000). TP53 evolution was examined in a variety of taxa, and sequence modifications were shown to be adaptive. The UPGMA rooted tree diagram of *H. sapiens* tumour protein p53 (TP53), transcript variant-1 mRNA sequence showed different clusters formation. Organism that originated from same ancestors having same e-value and 100% pair wise identity, are placed in same clusters whereas those which are distant from each other are placed in separate clusters. Phylogenetic reconstructions and an alignment of the *H. sapiens* tumor protein p53 (TP53) transcript variant-1 mRNA sequence were carried out using the "build" function of ETE3 3.1.3 (Huerta-Cepas et al., 2016) as it was implemented on the GenomeNet (<https://www.genome.jp/tools/ete/>). The multiple sequence alignment was supplied by the user.

The PhyML v20160115 model and settings were used to infer the ML tree: `-pinv e --alpha e --nclases 4 -o tlr -f m --bootstrap 1000` (Guindon et al., 2010). Out of 1000, branch supports are calculated.




Figure 2. Phylogenetic Analysis of Human Tp53 Gene

4. CONCLUSION

We may infer that the p53 protein plays a significant role as a tumour suppressor in bovines as well as other species, including humans. Animals that have a high proportion of identity and resemblance have more conserved amino acid sequences. Therefore, this work can offer a platform for researchers to determine animal comparative genomics

REFERENCES

- [1] Pintus, Sergey S., et al. "Phylogenetic analysis of the p53 and p63/p73 gene families." *In silico biology* 7.3 (2007): 319-332.
- [2] Pintus, S. S., et al. "Phylogenetic analysis of the p53 family." *Biophysics* 51 (2006): 571-580.
- [3] Sahoo, Pravas Ranjan, et al. "Phylogenetic analysis of p53 tumor suppressor gene of bos taurus through in silico platform." *Ovis* 92 (2018): 95.
- [4] Yang Z, Bielawski JP (2000). Statistical methods for detecting molecular adaptation. *Trends Ecol. Evol.* 15(12): 496-503.
- [5] Åberg, Emma, et al. "Evolution of the p53-MDM2 pathway." *BMC Evolutionary Biology* 17 (2017): 1-12.
- [6] Rutkowski, Rachael, Kay Hofmann, and Anton Gartner. "Phylogeny and function of the invertebrate p53 superfamily." *Cold Spring Harbor perspectives in biology* 2.7 (2010): a001131.

- [7] Fernandes, Andrew D., and William R. Atchley. "Biochemical and functional evidence of p53 homology is inconsistent with molecular phylogenetics for distant sequences." *Journal of molecular evolution* 67 (2008): 51-67.
- [8] Biscotti, Maria Assunta, et al. "The p53 gene family in vertebrates: evolutionary considerations." *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 332.6 (2019): 171-178.
- [9] Hiller, Thomas, et al. "Comparative analysis of 19 genital human papillomavirus types with regard to p53 degradation, immortalization, phylogeny, and epidemiologic risk classification." *Cancer Epidemiology Biomarkers & Prevention* 15.7 (2006): 1262-1267.
- [10] Jain AK, Allton K, Iacovino M, Mahen E, Milczarek RJ, Zwaka TP et al. p53 regulates cell cycle and microRNAs to promote differentiation of human embryonic stem cells. *PLoS Biology*. 2012; 10(2):e1001268
- [11] Woods YL, Lane DP. Exploiting the p53 pathway for cancer diagnosis and therapy. *Hematology Journal*. 2003; 4(4):233-247.
- [12] Shengkan J, Sebastian M, Woo SJ, Jennifer RW, Nebojsa M, Andrej S et al. Identification and characterization of a p53 homologue in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences*. 2000; 97(13):7301-7306
- [13] Soltis DE, Soltis PS. The Role of Phylogenetics in Comparative Genetics. *Plant Physiology*. 2003; 132:1790-1800.
- [14] Pintusa SS, Fomin ES, Ivanisenko VA, Kolchanov NA. Phylogenetic Analysis of the p53 Family. *Biophysics*. 2006; 51(4):571-580.
- [15] Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST and PSI BLAST a new generation of protein database search programme. *Nucleic acids research*. 1997; 25:3389-3402.

Citation: Akhil Kumar Pigili, Anil Kumar Babu Katika. *In Silico Phylogenetic Study of the Human Tp53 Gene: Insights into Evolutionary Dynamics*. *International Journal of Research Studies in Biosciences (IJRSB)*. 2024; 10(1):30-34. DOI: <http://dx.doi.org/10.20431/2349-0365.1001005>.

Copyright: © 2024 Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.